arXiv:1407.0191v1 [math-ph] 1 Jul 2014

Explicit energy expansion for general odd degree polynomial potentials

Asiri Nanayakkara* and Thilagarajah Mathanaranjan*†

*Institute of Fundamental Studies Hanthana Road, Kandy, Sri Lanka
†Department of Mathematics, University of Jaffna, Sri Lanka

(Dated: July 2, 2014)

In this paper we derive an almost explicit analytic formula for asymptotic eigenenergy expansion of arbitrary odd degree polynomial potentials of the form $V(x) = (ix)^{2N+1} + \beta_1 x^{2N} + \beta_2 x^{2N-1} + \cdots\cdots + \beta_{2N} x$ where β_k's are real or complex for $1 \leq k \leq 2N$. The formula can be used to find semiclassical analytic expressions for eigenenergies up to any order very efficiently. Each term of the expansion is given explicitly as a multinomial of the parameters β_1, β_2.... and β_{2N} of the potential. Unlike in the even degree polynomial case, the highest order term in the potential is pure imaginary and hence the system is non-Hermitian. Therefore all the integrations have been carried out along a contour enclosing two complex turning points which lies within a wedge in the complex plane. With the help of some examples we demonstrate the accuracy of the method for both real and complex eigenspectra.

PACS numbers: 03.65.-w, 03.65.Sq, 03.65.Ge

I. INTRODUCTION

A Hamiltonian with an odd degree polynomial potential

$$H = \frac{p^2}{2} + \beta_0 x^{2N+1} + \beta_1 x^{2N} + \beta_2 x^{2N-1} + \cdots\cdots + \beta_{2N} x \tag{1}$$

is $\mathcal{PT}$- symmetric when $\beta_0, \beta_2, ...\beta_{2N}$ are pure imaginary and $\beta_1, \beta_3, ...\beta_{2N-1}$ are real. It has real eigen spectra when $\mathcal{PT}$- symmetry is not spontaneously broken (i.e. when the wave functions are also $\mathcal{PT}$-symmetric). Moreover, the wave functions of H are usually required to vanish at infinity in various Stokes wedges to satisfy boundary conditions for quantization. Therefore, solving the Schrodinger equation directly to obtain eigen spectra for these systems is not a trivial task. Recently the eigenenergy spectra of general polynomial potentials have been investigated using spectral resolution method and extended WKB methods [1–12].

At present, the WKB theory is well developed and its methods are very important for many applications. The WKB method has been extended for obtaining higher order eigenenergies for potentials such as $V(x) = x^{2N}$[13]. For this system, the integrals in the each term of the expansion can be evaluated analytically in terms of Γ functions. Although, at the first sight, the problem of obtaining higher-order terms of WKB seems relatively simple for other systems, it has proved to be difficult due to singularities at the classical turning points [14]. However, the lowest order WKB method has been applied for obtaining eigenenergies of many $\mathcal{PT}$-symmetric potentials such as $V(x) = gx^2(ix)^\varepsilon$. Robnik et al. [14, 15] derived a simple formula for the semiclassical series for the potentials $V(x) = x^{2N}$ and obtained explicit formula for the WKB approximation of the eigenenergies for the same. The recurrence relations obtained by Robnik et al. are computationally much less time consuming compared to the WKB recurrence relations when the order increases. The reason behind this difference is that Robnik's recurrence relations involve only arithmetic operations

*Electronic address: asiri@ifs.ac.lk
†Electronic address: mathan@jfn.ac.lk

with rational numbers while WKB formulae involve operations of differentiation and collection of similar terms. In addition, they have derived almost explicit formula for the WKB terms for the energy eigenvalues of the potential $V(x) = x^{2N}$. However, use of the method developed by Robniks et al for general polynomial potentials are not possible due to the complicated nature of integrals involved.

Compared to the higher order WBK expansion, derivation of Asymptotic Energy Expansion (AEE) [16–19] for polynomial type potentials is relatively easy. All the integrals involved in Asymptotic Energy Expansion (AEE) of polynomial potentials can be evaluated analytically in terms of Γ functions. As a result, using Robnik's method in conjunction with AEE method, an almost explicit formula for semiclassical energies for even degree real polynomial potentials has been derived [6]. One of the important feature of this formula is that it contains parameters a_1, a_2, ...a_{2N-1} of the potential $V(x) = x^{2N} + a_1x^{2N-1} + a_2x^{2N-2} + \cdots\cdots + a_{2N-1}x$, explicitly. Since AEE is an expansion in reciprocal of energy, energy is also present explicitly in the formula. Recently, AEE method developed for even degree polynomial potentials has been applied for finding equivalent Hermitian Hamiltonians for non-Hermitian Hamiltonians successfully [20].

In a recent paper, Bender et al. [21] have developed a technique based on WKB theory to obtain the behavior of eigenenergy levels of the potentials of the type $V(x) = -igx^{2N+1}$ (for integer N) as g varies. The method is accurate enough to determine the critical points where the pairs of real eigenvalues get merged and become complex conjugate pairs. Due to the complicated nature of the integrals, this WKB method cannot also be applied for a general odd degree polynomial potentials. Recently, AEE method has been extended for the potential of the type $V(x) = \mu x^3 + ax^2 + bx$ [19]. Integrals in the AEE expansion of this system contain odd powers of $\sqrt{1-y^3}$, where y is the integration variable. Therefore, the integrals are evaluated with contours enclosing branch points 1 and ∞. However, the extension of AEE method to higher order odd degree polynomial potentials ($degree > 3$) failed to produce correct energy spectra. In this study it was found that the reason for this failure is the choice of branch points. In this paper we extend the AEE method to higher order odd degree polynomial potentials with correct choice of branch points for $degrees \geq 3$ and contours and derive an almost explicit analytic formula for asymptotic eigenenergy expansion for *arbitrary* odd degree polynomial potentials of the form

$$V(x) = (ix)^{2N+1} + \beta_1 x^{2N} + \beta_2 x^{2N-1} + \cdots\cdots + \beta_{2N} x \tag{2}$$

where N is a positive integer and $\beta_k \in \mathbb{C}$ for $1 \leq k \leq 2N$. The system has real eigenvalues when $V(x)$ is $\mathcal{PT}$- symmetric and eigen spectrum is complex otherwise. AEE expansion is valid for both real and complex asymptotic eigenenergies. The paper is organized as follows. In Sec. II we extend the AEE method for odd degree polynomial potentials with two terms and evaluate the accuracy of both real and complex eigenenergies. The main result of this paper, almost explicit analytic formula for asymptotic eigenenergy expansion for the potential (2) is derived in Sec. III. Two examples are given in section III to demonstrate the accuracy of the AEE method as well as the derived formula. In Sec. IV, a summary and concluding remarks are presented.

II. ANALYTIC SEMICLASSICAL ENERGY EXPANSIONS FOR $V(x) = (ix)^{2N+1} + bx$

In this section we investigate the two term odd degree polynomial potentials for the form

$$V(x) = (ix)^{2N+1} + bx, \tag{3}$$

where $b \in \mathbb{C}$. Consider the 1-D Schrodinger equation

$$-\hbar^2 \frac{\partial^2 U(x,E)}{\partial x^2} + V(x)\, U(x,E) = E U(x,E). \tag{4}$$

Substituting $P(x,E) = \frac{\hbar}{i}\frac{\partial U(x,E)/\partial x}{U(x,E)}$ in the above equation, we get

$$\frac{\hbar}{i}\frac{\partial P(x,E)}{\partial x} + P^2(x,E) = E - V(x). \tag{5}$$

Please note that $P(x,E)$ above corresponds to the derivative of the action in the usual WKB ansatz. The quantity $J(E)$ is now defined as

$$J(E) = \frac{1}{2\pi}\int_{\gamma} P(x,E)\, dx, \tag{6}$$

with the quantization condition $J(E) = n\hbar$. The contour γ encloses two turning points of $P_c = \sqrt{E - V(x)}$. For the potential in Eq.(3), Eq.(5) becomes

$$\frac{\hbar}{i}\frac{\partial P(x,E)}{\partial x} + P^2(x,E) = E - (ix)^{2N+1} - bx. \tag{7}$$

Let $\epsilon = E^{-\frac{1}{4N+2}}$ and $y = i\epsilon^2 x$. Then Eq. (7) becomes, after simplification,

$$\hbar\epsilon^{4N+4}\frac{\partial P(y,\epsilon)}{\partial y} + \epsilon^{4N+2}P^2(y,\epsilon) = 1 - y^{2N+1} - \frac{by}{i}\epsilon^{4N} \tag{8}$$

Now we expand $P(y,\epsilon)$ as a power series in ϵ,

$$P(y,\epsilon) = \epsilon^s \sum_{n=0}^{\infty} a_n(y)\,\epsilon^n \tag{9}$$

where s and $a_n(y)$ are determined below. Substituting (9) in (8) and equating coefficients of ϵ^0, we obtain $s = -(2N+1)$ and $a_0 = \sqrt{1 - y^{2N+1}}$ and Eq.(8) becomes

$$\hbar\sum_{n=0}^{\infty}\frac{da_n}{dy}\epsilon^{2N+n+3} + \sum_{i=0}^{\infty}\sum_{j=0}^{\infty} a_i a_j \epsilon^{i+j} = 1 - y^{2N+1} - \frac{by}{i}\epsilon^{4N} \tag{10}$$

and rearranging terms, we obtain

$$\left(\hbar\sum_{n=1}^{\infty}\frac{da_{n-2N-3}}{dy} + \sum_{n=1}^{\infty}\sum_{i=1}^{n-1} a_i a_{n-i} + 2a_0\sum_{n=0}^{\infty} a_n\right)\epsilon^n = 1 - y^{2N+1} - \frac{by}{i}\epsilon^{4N}. \tag{11}$$

Then coefficients a_n's are given by

$$a_n = \frac{-1}{2a_0}\left[\sum_{i=1}^{n-1} a_i a_{n-i} + \hbar\frac{da_{n-2N-3}}{dy} + \frac{by}{i}\delta_{4N,n}\right]. \tag{12}$$

In the above formula $a_n = 0\ \forall n < 0$. First four non zero a_n's for given N are

$$a_0 = \sqrt{1 - y^{2N+1}}$$

$$a_{2N+3} = -\frac{\hbar}{2a_0}\frac{da_0}{dy}$$

$$a_{4N} = -\frac{by}{2ia_0}$$

$$a_{4N+6} = -\frac{1}{2a_0}\left[a_{2N+3}^2 + \hbar\frac{da_{2N+3}}{dy}\right].$$

Now J can be written as

$$J(E) = \sum_{k=0}^{\infty} b_k E^{\frac{-(k-2N-3)}{4N+2}} \tag{13}$$

where

$$b_k = \frac{1}{2i\pi}\int_{\gamma} a_k(y)\, dy. \tag{14}$$

Now, in order to evaluate the integral (14), the contour γ is chosen such that it encloses the two branch points of $\sqrt{1 - y^{2N+1}}$ on the complex plane. There are 2N+1 branch points on the complex plane. The branch points which should be enclosed by the contour γ are $(-1)^N e^{iN\pi/(2N+1)}$ and $(-1)^{N+1} e^{i(N+1)\pi/(2N+1)}$. Note that these two branch points lie inside the Stokes wedges which are necessary for defining the above non-Hermitian problem correctly as an eigenvalue problem [7].

The integration is then carried out for each term and obtained the expression for $J(E)$ as

$$J(E) = b_0 E^{\frac{2N+3}{4N+2}} + b_{2N+3} + b_{4N} E^{-\frac{2N-3}{4N+2}} + b_{4N+6} E^{-\frac{2N+3}{4N+2}} \tag{15}$$

where,

$$b_0 = \frac{2\cos[\frac{\pi}{4N+2}]\Gamma[\frac{1}{2N+1}]}{\sqrt{\pi}\,(2N+3)\,\Gamma[\frac{1}{2} + \frac{1}{2N+1}]} \tag{16}$$

$$b_{2N+3} = -\frac{\hbar}{2} \tag{17}$$

$$b_{4N} = \frac{2bi\sin[\frac{\pi}{2N+1}]\Gamma[\frac{2}{2N+1}]}{\sqrt{\pi}\,(4N+2)\,\Gamma[\frac{1}{2} + \frac{2}{2N+1}]} \tag{18}$$

$$b_{4N+6} = \frac{2N\hbar^2\cos[\left(\frac{4N+1}{4N+2}\right)\pi]\Gamma[1 - \frac{1}{2N+1}]}{12\sqrt{\pi}\Gamma[\frac{1}{2} - \frac{1}{2N+1}]}. \tag{19}$$

By applying the quantization condition $J(E) = n\hbar$, $n = 0, 1, 2, ...$ the eigenenergies for $V(x) = (ix)^{2N+1} + bx$ can be obtained. Next we demonstrate the accuracy of the AEE method for odd degree polynomial systems with the help of two Hamiltonians. The first one is the $\mathcal{PT}$-symmetric Hamiltonian $H = p^2 + ix^5 + ix$ which possesses real eigen spectrum. The table I shows the first 12 eigenvalues of this system obtain with AEE in (15) as well as the numerical eigenenergies obtained with matrix diagonalization method. It is evident from the table I that AEE method for odd degree polynomial potentials produce accurate real eigenenergies even with four terms. As expected this method predict higher eigenenergies more accurately compared to the lower ones.

n	E_{AEE}	E_{Exact}
0	1.415221	1.624377
1	4.868558	4.820135
2	9.517600	9.522461
3	15.03904	15.03806
4	21.27666	21.27658
5	28.13384	28.13374
6	35.54327	35.54322
7	43.45471	43.45467
8	51.82880	51.82877
9	60.63369	60.63367
10	69.84293	69.84292
11	79.43411	79.43411

TABLE I: Comparison between calculated energy eigenvalues by AEE and E_{Exact} which is obtained by matrix diagonalization method for the Hamiltonian $H = p^2 + ix^5 + ix$. (where $\hbar = 1$)

The second illustration is the non-Hermitian non $\mathcal{PT}$-symmetric system given by $H = p^2 + ix^5 + (1 + i)x$. This system has complex eigen spectrum. The table II shows the first 12 eigenvalues of this system obtain with AEE in (15) as well as the numerical eigenenergies obtained with matrix diagonalization method. It is evident from the table II that AEE method for odd degree polynomial potentials can produce accurate complex eigenenergies even with four terms. Similar to the previous example, this method predict higher eigenenergies more accurately compared to the lower ones.

The analytic expression of $J(E)$ in (15) can also be utilized to investigate the asymptotic behavior of eigenvalues of potential in (3) analytically.

III. AEE OF GENERAL ODD DEGREE POLYNOMIAL POTENTIAL $V(x) = (ix)^{2N+1} + \beta_1 x^{2N} + \beta_2 x^{2N-1} + \cdots\cdots + \beta_{2N} x$.

In this section we use the same method used in Sec.II to obtain the AEE for the general odd degree polynomial potential in Eq. (2). The Eq. (7) now becomes

$$\frac{\hbar}{i}\frac{\partial P(x,E)}{\partial x} + P^2(x,E) = E - (ix)^{2N+1} - \sum_{k=1}^{2N} \beta_k \, x^{2N-k+1}. \tag{20}$$

n	E_{AEE}	E_{Exact}
0	1.385058 - 0.39235 i	1.529177 - 0.55265 i
1	4.857391 - 0.49947 i	4.826487 - 0.45524 i
2	9.511001 - 0.57092 i	9.514849 - 0.57341 i
3	15.03433 - 0.62534 i	15.03380 - 0.62425 i
4	21.27298 - 0.67002 i	21.27301 - 0.66976 i
5	28.13080 - 0.70831 i	28.13078 - 0.70813 i
6	35.54068 - 0.74204 i	35.54067 - 0.74193 i
7	43.45244 - 0.77233 i	43.45244 - 0.77226 i
8	51.82678 - 0.79992 i	51.82677 - 0.79987 i
9	60.63186 - 0.82532 i	60.63186 - 0.82528 i
10	69.84126 - 0.84890 i	69.84126 - 0.84887 i
11	79.43258 - 0.87096 i	79.43258 - 0.87093 i

TABLE II: Comparison between calculated energy eigenvalues by AEE and E_{Exact} which is obtained by matrix diagonalization method for the Hamiltonian $H = p^2 + ix^5 + (1+i)x$. (where $\hbar = 1$)

Let $\epsilon = E^{-\frac{1}{4N+2}}$ and $y = i\epsilon^2 x$. Then Eq. (20) becomes, after simplification,

$$\hbar\epsilon^{4N+4}\frac{\partial P\left(y,\epsilon\right)}{\partial y} + \epsilon^{4N+2}P^2\left(y,\epsilon\right) = 1 - y^{2N+1} - \sum_{k=1}^{2N}\beta_k\ \left(\frac{y}{i}\right)^{2N-k+1}\epsilon^{2k} \tag{21}$$

Now $P\left(y,\epsilon\right)$ is expanded as a power series in ϵ,

$$P\left(y,\epsilon\right) = \epsilon^s\sum_{n=0}^{\infty}a_n\left(y\right)\epsilon^n \tag{22}$$

where s and $a_n\left(y\right)$ are determined below. Substituting Eq. (22) in Eq. (21) and equating coefficients of ϵ^0, we obtain $s = -\left(2N+1\right)$ and $a_0 = \sqrt{1-y^{2N+1}}$ and Eq. (21) becomes

$$\hbar\sum_{n=0}^{\infty}\frac{da_n}{dy}\epsilon^{2N+n+3} + \sum_{i=0}^{\infty}\sum_{j=0}^{\infty}a_i a_j\epsilon^{i+j} = 1 - y^{2N+1} - \sum_{k=1}^{2N}\beta_k\ \left(\frac{y}{i}\right)^{2N-k+1}\epsilon^{2k} \tag{23}$$

and rearranging terms, we obtain

$$(\hbar\sum_{n=1}^{\infty}\frac{da_{n-2N-3}}{dy} + \sum_{n=1}^{\infty}\sum_{i=1}^{n-1}a_i a_{n-i} + 2a_0\sum_{n=0}^{\infty}a_n)\ \epsilon^n = 1 - y^{2N+1} - \sum_{k=1}^{2N}\beta_k\ \left(\frac{y}{i}\right)^{2N-k+1}\epsilon^{2k}. \tag{24}$$

Then coefficients a_n's are given by

$$a_n = \frac{-1}{2a_0}\left[\sum_{i=1}^{n-1}a_i a_{n-i} + \hbar\frac{da_{n-2N-3}}{dy} + \sum_{k=1}^{2N}\beta_k\ \left(\frac{y}{i}\right)^{2N-k+1}\delta_{2k,n}\right]. \tag{25}$$

In the above formula $a_n = 0\ \forall n < 0$. Now J can be written as

$$J\left(E\right) = \sum_{m=0}^{\infty}b_m E^{\frac{-(m-2N-3)}{4N+2}} \tag{26}$$

where

$$b_m = \frac{1}{2i\pi}\int_{\gamma} a_m(y)\,dy. \tag{27}$$

The contour γ encloses the two branch points of $\sqrt{1-y^{2N+1}}$ on the complex plane. The turning points and the contour used are the same as in the previous section. Note that these two branch points lie inside the Stokes wedges which are necessary for defining the above non-Hermitian problem correctly as an eigenvalue problem [7].

We studied explicit expressions for the polynomial potentials of order 3, 5, 7, and 9 and develop a general form for $a_m(y)$. Unlike in the even degree polynomial potential case, for odd degrees, there are two forms for the coefficient $a_m(y)$ in Eq. (27) depending on m is even or odd. Utilizing the procedure described in [18], we have obtain the general expression for $a_m(y)$:

For even m

$$a_{2m}(y) = -y^{\frac{2N-2m+1}{2}}\sum_{j=0}^{2m-1} A_{2m-j-1,j}\left[\frac{y^{(2N+1)/2}}{(1-y^{2N+1})^{1/2}}\right]^j \tag{28}$$

where

$$A_{s,l} = \frac{1}{2}\sum_{t=0}^{l-1}\sum_{i=0}^{s} A_{s-i,l-t-1}A_{i,t} - \frac{1}{4}\hbar i\,(s-2N(l+1)-2)\,A_{s-2N-2,l-1}$$

$$+\frac{1}{4}\hbar i\,(2N+1)(l-3)\,A_{s-2N,l-3} \tag{29}$$

and $A_{\alpha+\beta} = 0$ if $\alpha < 0, \beta < 0$ or $\alpha+\beta$ is even. $A_{2N,2} = \frac{(2N+1)i\hbar}{4}$ and $A_{4N-2k,1} = \frac{\beta_{2N-k+1}}{2}$ for $0 \le k \le 2N$.

For odd m

$$a_{2m+1}(y) = -\sum_{j=0}^{2m-1} A_{2m-j-1,,j}\left[\frac{y^N}{(1-y^{2N+1})^{1/2}}\right]^j; \qquad m \ge 4 \tag{30}$$

where

$$A_{s,l} = -\frac{1}{4}\hbar i\,(s+l(1-2N)-2N-1)\,A_{s-2N-1,l-1} + \frac{1}{2}\hbar i N\,(l-3)\,A_{s-2N+1,l-3} \tag{31}$$

and $A_{\alpha+\beta} = 0$ if $\alpha < 0, \beta < 0$ or $\alpha+\beta$ is even. $A_{2N-1,2} = \frac{iN\hbar}{2}$ and $A_{4N-2k-2,1} = \frac{\beta_{2N-k}}{2}$ for $0 \le k \le 2N-1$.

Next we obtain AEE coefficients b_n as

$$b_{2n} = -\frac{1}{2\pi i}\sum_{j=0}^{2n-1} A_{2n-j-1,j}\int_c \frac{y^{\frac{(2N+1)j}{2}+\frac{(2N+1)}{2}-n}}{(1-y^{2N+1})^{j/2}}dy \tag{32}$$

$$b_{2n+1} = 0$$

The contour integral in Eq. (27) can be evaluated in terms of Γ functions as

$$\int_c \frac{y^{\frac{(2N+1)j}{2}+\frac{(2N+1)}{2}-n}}{(1-y^{2N+1})^{j/2}}dy = \frac{4i^{\frac{2n+j+1}{2}}\cos[\left(\frac{2Nn+1}{4N+2}\right)\pi]\Gamma[1-\frac{j}{2}]\Gamma[\frac{3}{2}-\frac{n-1}{2N+1}+\frac{j}{2}]}{\left(1+\frac{(2N+1)j}{2}+\frac{(2N+1)}{2}-n\right)\Gamma[\frac{3}{2}-\frac{n-1}{2N+1}]}. \tag{33}$$

for odd j and when $\left(\frac{(2N+1)j}{2}+\frac{(2N+1)}{2}-n\right)$ is even. Otherwise, the integral vanishes. Then we have the expression for $J(E)$ as

$$J(E) = -\frac{\hbar}{2} + \sum_{n=0}^{\infty} d_n E^{\frac{-(2n-2N-3)}{4N+2}} \tag{34}$$

where

$$d_n = -\frac{2\cos[\left(\frac{2Nn+1}{4N+2}\right)\pi]}{\pi\Gamma[\frac{3}{2}-\frac{n-1}{2N+1}]}\sum_{j=0}^{2n-1} A_{2(n-j-1),2j+1} \times \frac{i^{n+j+1}\Gamma[\frac{1}{2}-j]\Gamma[2+j-\frac{n-1}{2N+1}]}{(1+(2N+1)(j+1)-n)} \tag{35}$$

$$d_0 = \frac{2\cos[\frac{\pi}{4N+2}]\Gamma[\frac{1}{2N+1}]}{\sqrt{\pi}(2N+3)\Gamma[\frac{1}{2}+\frac{1}{2N+1}]} \tag{36}$$

and $A_{2(n-j-1),2j+1}$ is given by (29) and (31). Note that parameters of the potential are now contained in coefficients $A_{2(n-j-1),2j+1}$ as multinomials in β_1,β_2, ... , β_{2N-1}.

Now, with following two examples, the accuracy of the formulas (34)-(36) is tested by calculating eigenvalues of two Hamiltonians using formulas (34)-(36) with the quantization condition $J(E) = n\hbar, n = 0, 1, 2, ...$ and comparing them with the exact eigenvalues obtained by numerical integration of the Schrodinger equation.

The first example is the Hamiltonian

$$H = p^2 + (ix)^7 + x^6 + ix^5 + x^2 \tag{37}$$

for this potential $N = 3$, $\beta_1 = 1, \beta_2 = i$, $\beta_5 = 1$ and $\beta_3 = \beta_4 = \beta_6 = 0$. First 25 non zero terms of the Asymptotic Energy Expansion (AEE) for this Hamiltonian were obtained from (34) - (36). The table III shows the AEE eigenvalues and the exact eigenvalues obtained by numerical integration of the Schrodinger equation for the Hamiltonian (37).

The second example is

$$H = p^2 + (ix)^9 + x^2 + ix \tag{38}$$

where $N = 4$, $\beta_8 = i, \beta_7 = 1$ and $\beta_k = 0$ for all $k < 7$. The AEE eigenvalues and the exact eigenvalues obtained by numerical integration of the Schrodinger equation for the Hamiltonian (38) are given in table IV.

It is evident from the above two examples that the formulas (34)-(36) derived for odd-degree general polynomial potentials produce very accurate eigenenergies for higher eigenstates. This is due to the fact that the large number of terms in the series could be now included in the calculation with the help of algebraic formulas (34)-(36).

n	E_{AEE}	E_{Exact}
0	1.5699863	1.4585541
1	5.1604902	5.1861926
2	10.477845	10.479973
3	17.144275	17.145466
4	24.969982	24.970596
5	33.833308	33.833555
6	43.646903	43.647038
7	54.343821	54.343906
8	65.870498	65.870553
9	78.182742	78.182781
10	91.243245	91.243274
11	105.01993	105.01995

TABLE III: Comparison between calculated energy eigenvalues by AEE and E_{Exact} which is obtained by numerical integration of the Schrodinger equation for the Hamiltonian $H = p^2+(ix)^7+x^6+ix^5+x^2$. (where $\hbar = 1$, $\beta_1 = 1, \beta_2 = i$, $\beta_5 = 1$ and $\beta_3 = \beta_4 = \beta_6 = 0$)

n	E_{AEE}	E_{Exact}
0	1.8453697	1.7229882
1	5.7301601	5.7860546
2	11.834124	11.847978
3	19.733814	19.732860
4	29.209838	29.209369
5	40.121601	40.121580
6	52.367271	52.367288
7	65.868043	65.868047
8	80.560282	80.560283
9	96.391051	96.391052
10	113.31531	113.31531

TABLE IV: Comparison between calculated energy eigenvalues by AEE and E_{Exact} which is obtained by numerical integration of the Schrodinger equation for the Hamiltonian $H = p^2 + (ix)^9 + x^2 + ix$. (where $\hbar = 1$, $\beta_8 = i, \beta_7 = 1$ and $\beta_k = 0$ for all $k < 7$)

IV. SUMMARY AND CONCLUDING REMARKS

We derived a simple formula for the semiclassical series for the general polynomial potential $V(x) = (ix)^{2N+1} + \beta_1 x^{2N} + \beta_2 x^{2N-1} \cdot \cdot \cdot \cdot \cdot + \beta_{2N} x$ using the recurrence relations obtained by AEE method. Almost explicit formula for asymptotic eigenenergy expansion is presented for the above potentials for any N. The formula can be used to find semiclassical analytic expressions for eigenenergies up to any order very efficiently. In a previous paper, similar expansions have been obtained for general even degree polynomial potentials. However, the Hamiltonian for odd degree polynomial potential considered in this paper is non-Hermitian and hence two branch points used for integration must lie inside the Stokes wedges which are needed for defining the non-Hermitian problems correctly as eigenvalue problems. It is important to identify that for such systems, the direct application of the WKB method to obtain higher-order terms in the expansion is found to be very complicated (if not impossible) due to the fact that integrals in the expansion coefficients

cannot be evaluated analytically. Although the WKB expansion and the AEE produce the same semiclassical series for the potential $V(x) = (ix)^{2N+1}$, they are completely different when the potential contains two or more terms. Therefore, the above explicit formula can be employed for obtaining semiclassical eigen spectra in the place of the higher order WKB method. With the aid of two examples we have shown the accuracy of the AEE method for both real and complex eigen spectra.

The AEE expansions are very useful in analyzing systems analytically. It can be utilized to find out how the level spacings, density of states and other quantities vary with energy and parameters of the potential. Recently AEE method found to very valuable in finding isospectral Hermitian and non-Hermitian pairs of Hamiltonians [20]. Since the odd degree polynomial potentials are non Hermitian and PT symmetric for certain combination of parameters $\beta_1, \beta_2 \cdots, \beta_{2N}$ the formulas derived in this paper will be valuable for finding equivalent Hermitian Hamiltonians for non-Hermitian Hamiltonians.

V. REFERENCES

[1] A. Voros, From exact-WKB towards singular quantum perturbation theory, Publ. RIMS, Kyoto Univ. 40,973 (2004).

[2] A. Voros, "Exact WKB integration" of polynomial 1D Schrödinger (or Sturm–Liouville) problem, Differential equations and the Stokes phenomenon (Proceedings, Groningen 2001), eds. B.L.J. Braaksma et al.,World Scientific, Singapore (2002), pp. 293.

[3] A. Voros, J. Phys. A32, 5993 (1999).

[4] A. Voros, J. Phys. A32, 1301 (1999).

[5] A. Voros, J. Phys. A 27, 4653 (1994).

[6] E. Delabaere, H. Dillinger and F. Pham, J. Math. Phys. 38, 6126 (1997).

[7] Kwang C. Shin, J. Phys. A38, 6147 (2005).

[8] Kwang C. Shin, Schrödinger type eigenvalue problems with polynomial potentials: Asymptotics of eigenvalues, arXiv:math/0411143.

[9] Kwang C. Shin, Half-Line non-self-adjoint Schrödinger operators with polynomial potentials: Asymptotics of eigenvalues, arXiv:math/0502522.

[10] Per Alexandersson, On eigenvalues of the Schrödinger operator with an even complex-valued polynomial potential, arXiv:1104.0593.

[11] Stefan Giller, J. Phys. A: Math. Theor. 41, 465202 (2008).

[12] G.A. Dobrovolsky, R.S. Tutik, J. Phys. A: Math. Gen. 33, 6593 (2000).

[13] C. M. Bender and S. A. Orszag, Advanced Mathematical Methods for Scientists and Engineers (Springer, 1999).

[14] M. Robnik, V. G. Romanovski, J. Phys. A:Math. Gen. 33, 5093 (2000).

[15] M. Robnik, V. G. Romanovski, Prog. Theor. Phys. Suppl. 139 (2000).

[16] A. Nanayakkara, Phys. Lett. A 289, 39 (2001).

[17] A. Nanayakkara and V. Bandara, Can. J. Phys 80, 959 (2002).

[18] A. Nanayakkara and I. Dassanayake, Phys Lett. A 294, 158 (2002).

[19] A. Nanayakkara, Can J Phys, 85, 1473 (2007).

[20] A. Nanayakkara and T. Mathanaranjan Phys. Rev. A 86, 022106 (2012).

[21] C. M. Bender, and Hugh F. Jones, Phys. Rev. A 85, 052118 (2012).